
Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.






The Home of Pacific Strains

MAR 5 1942 *

1942 CATALOG **CULTURAL DIRECTIONS**

VETTERLE & REINELT Hybridizing Gardens

CAPITOLA · CALIFORNIA



TO OUR PATRONS

Novelties... A great many of our visitors ask: "What have you that is new?" It is a difficult task to pull a fresh rabbit from a hat every time someone asks for it, and so it is with flowers. As we grow each year an entirely new crop of Begonias and Delphiniums, there are never two years alike.

In both Begonias and Delphiniums the advances in quality this year are far ahead of anything we have offered before. We do not deal in individual varieties but in strains which are constantly improved, so we do not offer

SANTA TO LOS BATOS CRUZ CAPITOLA ROAD топтевеч вяч MATSONVILLE HIGHWAY 41 ST. AVE VETTERLE REINELT SOQUEL CAPITOLA

just new varieties but new everything.

The only entirely new subject we are introducing to cultivation this year is the Pink Calla Lily listed on page 23. This is a species, the seed of which was imported from South America, and it's a real jewel for the garden.

Visitors are cordially invited to inspect our plantings during the season. The Polyanthus Primroses are at their best during March and the first part of April... In April and May the first crop of Delphinium seedlings is at its height ... Breeding stock of tuberous Begonias in our greenhouses is at its best in June and July ... In August and September over a million seedlings of tuberous Begonias can be seen in full bloom in our lath houses. After Labor Day the greenhouses are closed to the public, due to hybridizing.

How to Get There ... We are situated four miles out of Santa Cruz, as can be seen on the map. When coming from Santa Cruz, take the first side road to the right when past city limits and follow it to our nursery. If coming from Watsonville, turn left in Soquel, as indicated on map.

Join the AMERICAN BEGONIA SOCIETY

Descriptive bulletins on culture of all Begonias are issued monthly. Member-

ship, \$1 per year. Write to

AMERICAN BEGONIA SOCIETY 757 Athens Blvd. · Los Angeles, Calif.

BEGONIA TUBERS

The great majority of our tubers are produced annually from seed and ripen with us approximately the end of December. The size will vary anywhere from less than 1 in. in diameter to as much as 3 in. in diameter. The larger the bulb, the larger the plant one can expect. However, the size and quality of flowers are not governed by the size of the bulb.

For commercial distribution they are graded into three sizes, as shown in picture on page 4. Delivery, January to end of March.

SEEDLING TUBERS

- 1. Large Size; 2 in. up in diameter. This is the most preferred size, which produces large plants and if grown in pots requires minimum 8 in. or 9 in. containers.
- 2. Medium Size. $1\frac{1}{2}$ in. to 2 in. in diameter. Often this size is used for smaller pots, 7 in. in size, and will give excellent results for bedding.
- 3. Small Size. 1 to $1\frac{1}{2}$ in. diameter. This size will not give large plants the first season but it is satisfactory for bedding purposes or small pot plants. We do not offer these in our retail trade, as they require professional experience to bring best results.

CUTTING TUBERS

Nos. 4, 5 and 6. In some hanging varieties we offer tubers produced from cuttings, which differ only in form from the tubers produced from seedling plants. Cuttings often have to be grown for two years before they form sufficiently large tubers for sale. The cost of production and additional growing naturally raises the price of propagated varieties.

SPECIMEN TUBERS

No. 7. Each year we have a limited number of large tubers two to four years old, which have been grown in pots for breeding purposes. These, of course, are the finest Begonias we can offer for show purposes; to be grown in containers 12 in. in diameter.

CULTURE OF TUBEROUS BEGONIAS


Tubers... If early flowering is desired place tubers during January and February in a warm place in open trays. Moisten slightly once in a while until they come to life. As they begin sprouting, plant in flats in a mixture of peat and sand, leaf mold and sand, or peat alone, all of which are good media for developing a root system. Plant tubers 3 to 4 in. apart, so that they are ½ in: below the

surface. Keep uniformly moist but not too wet, in a warm place, well lighted, until 3 or 4 in. of growth develops. Then plant in a permanent location or in pots, as desired. The front of the plant is always where the tips of the leaves are pointing. Often if dormant tubers are planted in open ground, especially if it is too cold or wet, a number of them may rot and the planting will be uneven, with some coming earlier and some later, with the plants facing in haphazard ways. Started first in flats, they will develop a splendid root system which is necessary for the forming of fine specimen plants.

Transplanting . . . Before the plants become too large and crowded in flats, transplant in open ground or pots, taking care not to disturb the root system too much. Outdoors any light soil containing a lot of humus will grow good begonias. Heavy soils should have a strong application of a well-rotted manure, leaf mold or sand to lighten them. Light, gravelly or sandy soils will benefit greatly by the application of peat, leaf mold or well-rotted manure. If planting in pots, soil should be much lighter than that used outdoors; two-thirds coarse leaf mold and one-third sand or sandy loam will give excellent results.

Feeding . . . If well-grown specimens are desired, additional feeding will be required, either with fish or cottonseed meals, both of which give fine results. The best method of using these fertilizers is to mix them with the soil that goes into the lower half of the pot, so that the roots will gradually reach into this area. If planting outdoors, a heaping tablespoonful for small plants and two for large plants will be sufficient when placed in the lower half of the planting hole. In pots a tablespoonful for a 6-in. pot or a small handful for an 8-in. pot will be enough to carry the plants through the season. See that the fertilizer does not come into contact with the stem or leaves of the plant, as both form a mold in the early stage of fermenting which would attack the growing tissues and destroy the plant.

Watering . . . After transplanting, gentle overhead watering is best until the plants are well established. Too heavy watering during the young stage may pack the soil, keeping it too wet, thereby causing the soil to go sour. Later, when the plants are in full growth, they will require a steady supply of moisture as the root system is shallow and any drying out will give a set-back to normal development.



TUBEROUS BEGONIA PLANTS

Seedling Plants . . . Commercially all tuberous begonias are produced annually from seed. Seed sown during January and February will produce flowering plants, from July on, and a tuber upon maturing in December. From then on the tuber, each year, grows in size forming a larger and larger plant. For the benefit of those who have no facilities or time for starting tubers, or for growing begonias from seed, we offer seedling plants of all types and colors listed on pages 7, 8, 9, 10, 11 and 12.

These seedlings are twice transplanted and hardened off before shipping so that upon arrival they can be planted directly outdoors in a permanent location. The shipping season begins May 15 and ends June 15. Later plantings would not bring sufficiently large plants to give enough flowers the same season. Plants that are planted in May and early June will begin to bloom by the end of July and reach their height of flowering season during August-September and continuing on into November.

Regions with warm summers and long cool autumns, such as most parts of California, will get better results from seedlings than from tubers. Tubers start blooming earlier and the flowers often burn during the hot summer months. If one desires a long season of flowering, tubers can be planted for early bloom and seedlings for continuing the season until the end of autumn. For bedding purposes seedlings are preferable as they grow uniformly and form a mass of color.

Seedlings are shipped successfully to all parts of the United States but we guarantee safe delivery only on the Pacific Coast and cannot be responsible for shipments east of the Rocky Mountains.

CULTURE OF TUBEROUS BEGONIAS

Seed... Sow from January to March. Place one or two inches of gravel in flats or seed pans, to insure good drainage, over which place a fairly coarse mixture of two-thirds leaf mold and one-third peat, about one inch deep. Smooth the surface with the same mixture, finely sifted, not more than one-eighth inch deep but do not press it down. It is necessary for the surface to be of a spongy character, so that when the seed germinates, the young roots can get into it. If surface is too fine and packed, the seedlings will often fall over, as the young roots are unable to get in . . . Place the pans in shallow water until thoroughly soaked up from below, then broad-

cast the seed. Cover with glass and a sheet of paper or keep it in the dark until germination takes place. Night temperature of 65 to 75 degrees Fahrenheit is necessary for quick germination. Lower temperature than 65 degrees will considerably slow up germination, with poorer results . . . As soon as germination takes place, take the paper off and in three or four days lift the glass also, otherwise the little seedlings will get too spindly. Warm temperature, protection from direct sunlight, and uniform moisture, are absolutely necessary. Even a slight drying out of the surface will be fatal to the delicate young plants. Very gentle overhead watering should be used when necessary . . . Dampening off will occur only if flats are kept too wet in greenhouse without proper ventilation. Watering with a weak solution of Clorox, about 2 to 4 per cent, we found as best cure and preventive.

Transplanting . . . When the third leaves are developed, transplant one inch apart in flats containing the same mixture as for sowing. No finely-sifted surface is necessary any more. Do not put more than one to one and one-half inches deep of soil in your flats. Deep flats filled with several inches of soil are not necessary. Usually they do not drain well, soil will get sour, and checking of the growth will be the result. Before they get too crowded, transplant in the same manner again farther apart, until strong enough to be planted out in the open or, if desired, potted up. If planting in open ground, enrich the soil liberally with well-rotted cow or sheep manure and a sprinkling of bone meal. If soil is heavy, add plenty of either leaf mold, peat or sand or a mixture of all, so that the ground will be light and porous. Plant in a shaded position, such as under the trees or north side of the house, where direct sunlight cannot reach them. Keep well watered; fine, overhead sprinkling preferred.

Digging and Storage . . . In autumn, when the foliage turns yellow, withdraw the water gradually and when all growth dies down entirely, take out, wash off all soil, taking care not to bruise the tubers, dry in sunlight for a day or two until thoroughly dry; then store in open flats in cool, dry place. See that all particles of the old stem are removed until healthy tissue shows; otherwise, if left on, they will decay and destroy the tuber.

SINGLE TYPES OF TUBEROUS BEGONIAS



GIANT SINGLE



SINGLE FRILLED



SINGLE CRESTED

TUBEROUS BEGONIAS

Giant Single Type . . . The individual flowers of this group often reach astonishing size of from 8 to 12 inches in diameter; carried on strong, stiff stems. They are very decorative as large specimen plants in tubs or for bedding purposes, where they will form a solid sheet of color.

COLORS

White	Salmon	Orange
Pink	Yellow	Scarlet
Rose		Dark Red

Single Frilled Type (Crispa)...Perhaps the most adorable of the single types. The large, individual flowers are beautifully frilled and ruffled at the edge.

COLORS

Solid Colors-	-White	Yellow	Scarlet
	Pink	Orange	e Dark Red
Variations-	-Rose		Light Salmon
	Crimson	Rose	Dark Salmon
	Apricot		

Fascination —On cream background, edged in rosy red.

Single Crested Type (Cristata)...The accompanying picture speaks for itself as to the beauty of this type, which we have highly improved in recent years. Dwarf, stocky growth, with masses of flowers, make this type suitable both for bedding and show purposes.

COLORS

White	Salmon Shades	Orange
Pink Shades	Yellow	Scarlet
Rose Shades		Dark Red

PRICE OF SINGLE TYPES

Tubers... Delivery, January to March.

Large size . . . 20c each; \$2.00 per dozen;

\$14.00 per 100.

Medium size . . . 15c each; \$1.50 per dozen; \$10.00 per 100.

Seedling Plants... Delivery, May-June. \$1.00 per dozen; \$7.50 per 100. F.O.B. Capitola. Seed... 50c per packet; 25c per half packet.

THE MOST IMPORTANT FACTORS FOR GROWING TUBEROUS BEGONIAS

The species used for the development of tuberous Begonias grown today came originally from South America, chiefly Bolivia and Peru, although in a tropic region they are found in their native state only at higher altitudes. This means that they require a cool climate for best development. In regions with extreme heat and humidity during summer they will not perform well, if at all. They are not suitable for house plants.

The Begonias are strictly summer flowering plants, dormant during December and January. Therefore, they cannot be grown in winter in states like Florida, etc. Well-matured tubers can be forced under glass but even then the earliest flowers will appear only by the end of April or early May. They always have to be protected against strong sunshine and perform best in semi-shade. Curled and shiny foliage is a sign of too much sun; too dense shade will produce spindly growth without flowers. They require constant moisture, as their root system is very shallow; and also perfect drainage.

Failures will occur where any of the cardinal points are disregarded. Falling off of buds before they develop will happen either if the plants are not in a healthy state or during hot weather. Small pots that dry out very rapidly or, again, too large pots which stay constantly wet, will cause it also. Overfeeding is the most common mistake, resulting in soft, glassy texture of the foliage, curling under of leaves, gradually wilting and dying off. Overwatering, chiefly in pots with poor drainage, will often have the same results.

Pests are practically non-existent outdoors. Occasionally greenhouses where a variety of plants are grown will have Begonias attacked by *Cyclamen mite* in some districts. Brown, rusty streaks on the foliage and buds are the first sign of this. Spraying with light oil sprays such as Volck, and isolating infected plants, are the best checks.



DOUBLE CAMELLIA TYPE

The largest and most popular of the double types. Individual flowers, from 5 to 8 in. in diameter, resemble camellias and roses in all forms and variations of color.

COLORS

Solid, Uniform Colors, with slight variation only:

White Rose American Beauty Cardinal Red Dark Red Orange
Pink Crimson Rose Salmon Red Scarlet Yellow Flame Orange

Salmon Shades, varying more or less within themselves:

Salmon Rose Shades Blush and Pink Salmons
Apricot and Orange Salmons Yellow and Apricot Salmons

PRICE

Tubers... Delivery, January-March. Large size . . . 25c each; \$2.50 per dozen; \$18.00 per 100 Medium size . . . 20c each; \$2.00 per dozen; \$14.00 per 100

Seedling Plants . . . Delivery, May-June. \$1.50 per dozen; \$10.00 per 100. F.O.B. Capitola. Seed . . . \$1.00 per packet; 50c per half packet.



DOUBLE CAMELLIA PICOTEE SHADES

We developed this group by inter-crossing the large camellia type with the small marmarota type. The flowers are slightly smaller in some instances but of beautiful camellia form, with two-toned combinations of color with a definitely contrasting edge.

COLORS

Red Shades

Pink-Rose Shades

Salmon Shades

PRICE

Tubers . . . Delivery, January to end of March.

Large size . . . 25c each; \$2.50 per dozen; \$18.00 per 100 Medium size . . . 20c each; \$2.00 per dozen; \$14.00 per 100

Seedling Plants . . . Delivery, May-June. \$1.50 per dozen; \$10.00 per 100. F.O.B. Capitola. Seed . . . \$1.00 per packet; 50c per half packet.



DOUBLE ROSEBUD TYPE

Flowers of the most exquisite form, resembling perfect rosebuds, with beautiful, two-tone, pastel-shade coloring. Very tall, sturdy growth makes it especially desirable for growing large specimen plants in tubs for decorative purposes.

COLORS

Pastel Shades in Blush Pink Rose Salmon Flame Salmon

PRICE

Tubers... Delivery, January-March. Large size . . . 25c each; \$2.50 per dozen; \$18.00 per 100 Medium size . . . 20c each; \$2.00 per dozen; \$14.00 per 100 Seedling Plants . . . Delivery, May-June. \$1.50 per dozen; \$10.00 per 100. F.O.B. Capitola.

Seed . . . \$1.00 per packet; 50c per half packet.



FIMBRIATA PLENA TYPE

Double frilled, or often called Carnation type. This group has been enormously improved in the last few years, the flowers in some cases reaching the size of the Camellia type, with more and more refined form. Its strong, bushy growth makes it exceptionally desirable for bedding and pot plants.

COLORS

White Rose Salmon Red Orange
Blush Crimson Rose Yellow Scarlet
Pink Salmon Apricot Dark Red

PRIC

Tubers... Delivery, January-March. Large size . . . 25c each; \$2.50 per dozen; \$18.00 per 100

Medium size... 20c each; \$2.00 per dozen; \$14.00 per 100

Seedling Plants... Delivery, May. \$1.50 per dozen; \$10.00 per 100. F.O.B. Capitola.

Seed . . . \$1.00 per packet; 50c per half packet.



DOUBLE HANGING TYPE

Its hanging habit, with great masses of blooms, makes them favored for the decoration of greenhouses, open verandas or sun porches, where they can be protected against strong winds and light. We have been improving this type by interbreeding them for a number of years with the double Camellia type for the size of flower and variety of color. Our this year's offerings are both further advanced, combining medium-sized flowers and good hanging habit.

COLORS

White	Pink	Crimson Rose	Yellow	Scarlet
Shell Pink	Rose	Salmon	Orange	

PRICE

Tubers... Delivery, January-March. Large size . . . 25c each; \$2.50 per dozen; \$18.00 per 100 Medium size . . . 20c each; \$2.00 per dozen; \$14.00 per 100

Seedling Plants . . . \$1.50 per dozen; \$10.00 per 100. F.O.B. Capitola.

Seed... This year we can offer only a small quantity in the complete mixture of the above and all new crosses made this season. Price, \$1.00 per packet; 50c per half packet.



NAMED VARIETIES

We have been selecting and propagating a number of seedlings of the basket Begonias which are outstanding in color, form of flowers and hanging habit. All are propagated by cuttings, forming rather irregular tubers in comparison to the seedling tubers which are smooth and round. It takes several years to propagate a sufficient amount of tubers for introduction and we are so far able to offer only the two following varieties:

Pink Shower . . . The individual flowers are a delicate cerise-pink, holding the beautiful form to the very end of the season in such profusion that the plant portrays the name it bears exactly.

Avalanche . . . Recent introduction and a good companion to Pink Shower, as it is almost a white counterpart of it. A slight greenish tinge in the center of the flowers gives it added attraction.

Price ... 75c each; \$7.50 per dozen. No seed of the above-named varieties is available, as they are produced only from cuttings.

CULTURE OF HANGING BEGONIAS

Hanging basket Begonias have the same cultural requirements as all other types, with minor exceptions. For good results one should have large tubers, as the larger tuber will have more shoots come from it and consequently the plant will be larger and more effective. Tubers can be started from January to March, the same as any other type, but when the growth reaches three or four inches transplant them in a light mixture of soil; if possible, two-thirds coarse leaf mold and onethird sand, as they are very sensitive to perfect drainage. Containers should be large enough; a minimum of 8 inches in diameter for small tubers and up to 12 inches for large ones. Shallow pots, if possible glazed, are better than wire or any other type basket, as the plants do not dry out in these types so severely and can produce far better growth. Wire baskets, lined with moss, can be utilized but one can not expect to grow excellent specimens by this method. Tubers started in peat will develop a sufficiently large root system so that they can be planted immediately in pots large enough to carry them through the season.

Feeding . . . It will be necessary to mix a small handful of fish meal with the soil going into the lower half of the pot, which will supply the plant with sufficient nutrients to start with. Later in summer, when the plant shows a decline in growth, another small handful dug into a shallow trench around the edge of the pot and covered with soil will revive it completely, and bring a new profusion of blooms.

Pinching . . . Some types, which do not show more than one or two shoots at the beginning of the season, should have the heart pinched out when the growth reaches the first flower bud. This will induce the side shoots to develop fully and form a better balanced plant.

VISIT OUR GARDENS

in July and August
when the Basket Begonias are
at their best



CHOICE SELECT VARIETIES OF TUBEROUS BEGONIAS

Contrary to the practice of propagating named varieties of tuberous Begonias, we offer selected seedling tubers instead. Propagation of individual varieties is costly and slow, so that by the time a sufficient amount has been reproduced for introduction they are already obsolete as compared with seedlings. Breeding is advancing so rapidly that today's choice is the discard of tomorrow. During flowering season we select and mark all the choicest new seedlings which are above average in size, form and color. We are offering tubers of these to connoisseurs who desire only the finest for their gardens. For show purposes and greenhouse decoration they are the last word in tuberous Begonias. In this group we supply all the double types and colors listed on the previous pages, besides including also all the new color developments derived from our crosses preliminary to introduction. "Apple Blossom," a double Camellia, is supplied in this group only.

This year's advance in breeding brought results far beyond expectations. Some colors, such as Cardinal Red, Red Salmon, Rose and Salmon Rose tones, have reached extremely large size. The average quality of all other colors is above anything we have previously offered.

In this group we also have a limited supply of specimen tubers, two or more years of age, which have been grown for breeding purposes in our greenhouse displays. When ordering these please specify second and third choices of color, as we never have a large quantity of any type or color in stock.

PRICE OF CHOICE SELECT VARIETIES Tubers Delivered January to March

Large size . . . 50c each; \$5.00 per dozen; \$37.50 per 100 Medium size . . . 40c each; \$4.00 per dozen; \$30.00 per 100 Specimen tubers . . . \$1.00 each; \$10.00 per dozen

SPECIAL HINTS

To grow Begonias to their utmost perfection requires, of course, the finest quality of tubers. Size does not matter; smaller tubers of a fine variety will produce as large a flower as the larger tubers, but the latter will form larger plants with more flowers. To gain on flower size, break out the weaker sprouts when tubers come to life—leaving only one of the strongest which will then develop into a very strong stalk bearing larger flowers than if several were left. This does not apply to the hanging type, where as many shoots as possible are wanted for good specimens.

Careful growing and feeding, as described on previous pages, should be followed. Disbudding of female flowers is of no help whatsoever. Very large flowers, until the end of the season, can be had by cutting out the sideshoots when young, so that all strength goes into the single stalk. This can not be performed on all varieties; only where one can cut out the young offshoot in the axil without bruising the tissue of the stem. Dusting wounds with charcoal or exposing them to sunlight for a time will heal them perfectly.

If large tubers, which form more shoots, are available, these can be left until 3 to 4 inches high. Then cut the weaker ones at the base, so they contain the basal ring, and then root them in sand in a closed propagation bed, either with or without bottom heat. They will form roots in 3-6 weeks and can then be potted and will bloom nicely the same season, although only a small tuber will be formed. In this manner extra choice plants can be propagated. The cutting tubers in their second year do not branch as heavily as seedling tubers and consequently bring the largest and most perfect blossoms. One can preserve fine varieties by this method and gradually accumulate a collection of perfect plants.



PACIFIC STRAIN OF

Although a comparatively short time has passed since our strain of Delphinium was introduced to the gardening world, the enthusiastic acclaim of amateurs and professionals alike has made us feel that we have contributed a step in the progress of better Delphiniums. At 1939 Oakland Spring Garden Show, which is the annual flower event of the Pacific Coast, our Delphiniums received the gold medal for the most outstanding display in the show, in open competition with all other spring flowers. These plants were field-grown seedlings in their maiden bloom, dug and balled the day before the show, without being forced in any way. We were again awarded the gold medal in 1940, but were unable to show in 1941 because of rainy weather which ruined all our flowers.

Contrary to the popular practice of naming individual plants, our aim is to produce a group of individual plants resembling each other as closely as possible, keep interbreeding them until the progeny brings a high percentage of specimens true to description, and then introduce it as a named series. In this manner we can offer new progressions of each series each year as the progeny always supersedes the parents. The names of our Delphinium series, with the exception of the blues, which have descriptive names, have been chosen from Tennyson's "Idylls of the King."

RECENT INTRODUCTIONS

Galahad Series . . . Our greatest achievement in giant whites, fully worthy of its name, which possesses all the qualities one expects to find in a fine Delphinium. The standard attained in this series is perhaps responsible for the wide popularity it met, ranking in demand next to the blues. The clear white flowers, with white bees, are of large size, reaching often 3 inches in diameter, combined with glistening, heavy texture. The 1942 series are outstanding in size, texture and purity of color, which we only dreamed of when the first series of Galahad was introduced.

Blue Bird Series ... A true blue Delphinium has been the hardest thing to produce and we have been reserving the name Blue Bird to signify our best achievement in this color. We can safely recommend this series as being the clearest medium blues, with white bees. They not only come true to color but carry well-formed, round flowers $2\frac{1}{2}$ inches in diameter, on very long, graceful spikes. The habit is all one could desire in modern Delphinium and this series sets a new standard in blues of today.

Cross-Pollinated Seed . . . Price \$3.00 per packet; \$1.50 per half packet.

DELPHINIUM HYBRIDS

The qualities we look for in Delphiniums are not only huge flowers and spikes, regardless of other characteristics, but on the contrary we often use plants of smaller size if they have outstanding qualities in color or form. Our aim is to produce large flowers properly proportioned to the size of spikes, so as to maintain balance and retain the grace of the entire plant. Another important feature in our developments is comparative mildew-resistance. Over ten acres of seedlings are being grown this year for selection, from which only the finest members combining all the desirable characteristics are retained for seed production. This enables us to produce very high quality seed, giving customers the advantage of the newest developments.

NAMED SERIES

Cameliard Series . . . A clear lavender self with white bee, possessing classical perfection of form of the individual flowers, which are very round and attain very large size, averaging 3 inches in diameter. Beautifully balanced spikes, combined with excellent growing habits.

Guinevere Series... A clear, pink-lavender self, with white bee; with very large individual flowers averaging up to 3 inches in diameter. This is a true exhibition type and a beautiful garden plant.

King Arthur Series... This is still one of the most brilliant Delphiniums we have so far developed. The color is a rich, royal purple, with a velvety texture and large white bee, and the beautifully formed, long spikes are carried on thin, woody stems.

Black Knight Series... The darkest violet. Individual flowers $2\frac{1}{2}$ to 3 inches in diameter; of beautiful round form, with heavy velvety texture, which gives the color luminosity and vividness not seen before in Delphinium. Very long, well-formed spikes: a true show flower with black bees.

Summer Skies Series . . . As the name indicates, this is the blue of a summer sky, with white bees representing fleecy clouds; a color long sought in the hybrid Delphiniums.

Blue Jay Series... Every Delphinium lover seeks the true blue color, which is so rare in nature and so difficult to obtain. These series are our bid toward attaining it. The color is striking, clear medium to dark blue; very intense and alive, with dark, contrasting bee.

Round Table Series... As the name implies, this represents all of the color combinations of the Knights and their Ladies which will eventually be introduced. This year some three hundred different crosses were used to make up this group, combining all the newest color developments and possessing size that will make them invaluable for show purposes.

Cross-Pollinated Seed... Price \$2.50 per packet; \$1.75 per half packet. Collection of five packets, \$10.00; collection of five half packets, \$5.00.





PACIFIC STRAIN OF

STANDARD COLORS

Besides the named series we are offering several groups which are not yet segregated to such a degree that they would come true to color. They are the best we have been able so far to produce in their respective colors and are fully equal in character of growth and size of flowers to the named series. Produced by artificial cross-pollination to intensify the desirable characteristics in the highest degree. It is the same seed we use for our own planting, and by releasing it to the public we feel that we are giving them an equal chance. The percentage of show specimens from this seed is the highest obtainable.

Lavender Pastel Shades . . . Lavenders with smoky pink in the petals and dark brown or black bee; highly mildewresistant.

Lilac Shades . . . Pure lilac tones, white bee predominant.

Light Violet Shades . . . Flowers very large, ranging from light violet to lilac tones.

Light Blue Shades . . . Ranging from light to medium blue, with white to fawn bee.

Medium Blue Shades . . . Clearest blues are in this group, with white and dark bees.

Dark Blue and Blue Violet Shades . . . Richest coloring in delphinium comprises this group. A combination of dark blue and violet is further intensified with dark bees.

Clear White... These are slightly different from the Galahad Series, the flowers not being as large; however, they are the purest white, with velvety texture. Very heavy branching and long trusses of flowers make them ideal for cut flower purposes.

White, Black Bee . . . The best whites with dark bees we have been able so far to produce.

Cross-Pollinated Seed... Price \$2.00 per packet; \$1.00 per half packet. Collection of four colors, \$7.00; half packets, \$3.50. Collection of eight, \$13.00; half packets, \$6.50.

JOIN AMERICAN DELPHINIUM SOCIETY

Beautiful Yearbook containing history of the newest developments in the Delphinium world. Published annually. Membership, \$2.00 per year. Write to

THE AMERICAN DELPHINIUM SOCIETY
Hinsdale, Illinois

DELPHINIUM HYBRIDS

Delphinium Plants . . . From March to the end of June and from September to the end of November we are offering strong, young plants, in twelve assorted shades, as marked for cross-pollinized seed. Transplanted, they will grow rapidly on and bring far better results in a shorter time than the old clumps. We prefer distribution of young seedlings rather than the one- or two-year-old clumps. which very seldom bring good results, as they are quite exhausted and it takes a year at best to get them re-established, during which time the flowers are not at their best. Young seedlings will produce show specimens within four or five months after planting and continue blooming for several years. In California interior valleys where the climate is too warm, autumn planting, September and October, is preferable. Plants having time to establish good root systems during winter will bring beautiful spikes in early spring. Cooler regions will give good results from both autumn and spring planting, except in heavy, cold soils, where spring planting is preferable.

Seedling Plants... \$2.00 per dozen; \$15.00 per 100. F.O.B. Capitola.

Self-Pollinated Seed... Besides the cross-pollinated seed, which is produced only in small quantities, we also supply self-pollinated seed to cover the commercial trade. Of the one hundred thousand seedlings which we grow annually, 40 to 60 per cent are rogued out, leaving only the best for the production of self-pollenized seed which we list below.

Named Series . . . For full description, see page 17.

Galahad Series Blue Bird Series

Seed . . . Price per packet, \$1.50; per half packet, 75c.

Cameliard Series Guinevere Series
King Arthur Series Black Knight Series
Summer Skies Series Blue Jay Series
Round Table Series

Seed . . . Price per packet, \$1.25; per half packet, 65c.

Standard Colors . . . For description, see page 18.

Lavender Pastels Light Blues
Lilac Shades Medium Blues

Light Violet Shades Dark Blues and Blue-Violet
Clear White White with Black Bee

Seed . . . Price per packet, \$1.00; per half packet, 50c. Collection of eight, \$7.00; half packets, \$3.50.





THE CULTURE OF OUR

Seed . . . Delphiniums can easily be grown from seed which can be sown practically any time of the year, according to the climatic conditions, equipment available or time of flowering desired. Under California conditions, for early spring flowers, sow from June to September; for midsummer blooms, December to January; and for fall blooms, February to April. In regions with severe winters, the early summer sowing, from June to July, is recommended, so that the little plants are well established before the winter sets in. Otherwise, the most practical time to sow is early spring. Seed can be kept in good condition for several years if kept in air-tight containers in a refrigerator. In fact, it is practicable even with fresh seed to place it between two moist blotting papers and leave it directly under the freezing compartment of the refrigerator for at least a week to induce higher germination. For best results, sow in flats in a mixture of two-thirds coarse leaf mold and one-third loam, covering slightly with the same mixture, and moisten thoroughly. To prevent evaporation, cover the flats with newspaper and glass until germination takes place. Immediately after the young plants begin to appear, both the glass and the newspaper should be taken off; however, the plants should be kept shaded and kept moist constantly.

Germination... If the seeds do not germinate 100 per cent and the flats are exposed to strong light immediately after the first few

plants have appeared, the rest may not germinate at all. Artificial heat can be used for germination in early spring and it will give far better results than the natural heat in summer. To get the best results in germination, bottom heat is necessary, with a cool temperature overhead. This is well supplied under glass in early spring; however, in summer, due to hot weather conditions, it is usually the opposite and this is why many people have failures even with the very freshest seed. At temperatures of 85 degrees Fahrenheit and up, the germination is often very poor and what germinates stands a very good chance of being simply cooked. To prevent this, after the seeds are sown in flats, place them on the floor in a cool room or shed, where they can be kept dark for the first ten days, until germination takes place. Then give light and fresh air, but keep them well protected against any drying out. We have repeatedly checked on seeds which were sown under greenhouse conditions in summer or the cool shed method or a well-shaded lath house, and the results in each case were from 20 to 50 per cent better under lath house conditions than under the greenhouse conditions. Under slow, cool germination practically every seed will gradually come up. Under quick, hot conditions only a few will germinate, the rest remaining dormant in the soil. Dampening off will occur only if flats are kept too wet in the greenhouse, without proper ventilation.



DELPHINIUM HYBRIDS

Transplanting . . . When the second leaves are developed and before the plants are too crowded, prick in flats 3 inches apart in a mixture of two-thirds sandy loam and onethird leaf mold. Keep shaded for two or three weeks; then gradually give more light and, when larger, harden off in full sunlight before planting out in permanent position about two or three feet apart. Open, sunny location is necessary for best development. In shade they will grow too spindly, with only small flower spikes. If too close to walls, they will mildew more than in the open. The ground should be well prepared for planting. A liberal application of well-rotted cow or sheep manure, with a sprinkling of bone meal mixed with the soil, which should be dug a foot deep, will produce fine growth. Heavy, wet soils will require a slight addition of lime. Good drainage and uniform supply of moisture during the growth is essential.

Feeding... When the first crop of blooms has faded, cut the flower spikes off just above the foliage and keep slightly dry for two or three weeks, to give the plants time to rest before the new shoots appear above the ground. When this takes place, cut the rest of the old stock off, sprinkle a teaspoonful of ammonium phosphate around each plant, rake it into the soil slightly and water thoroughly. From the new shoots appearing from the ground select two or three of the strongest and break the rest out. The remaining ones will develop into fine spikes again. Do not force a new growth

late in autumn; rather, keep the plants on the dry side, because if forced into bringing a third crop late in the season the plants will soon exhaust themselves and gradually die.

Diseases . . . MILDEW will attack plants grown close to a wall, or planted thickly together, especially later in the fall. Sulfur, dusted on the foliage, will act as a preventative before the disease is established. GREEN FLOWER-So far this virus disease has been reported only west of the Rocky Mountains and is most prevalent close to ocean areas. It is transmitted by a species of leaf hopper from infected weeds to the Delphinium plants. All plants showing signs of the disease should be immediately discarded. CROWN ROT may be caused by several fungi prevalent in the eastern and southern sections of the United States where high temperatures and humidity prevail during the growing season. For best information send parts of infected plants to your state university for identification. Further information on this subject can be found in the Yearbooks of The American Delphinium Society.

Length of Life . . . Length of life of Delphinium plants is governed by several factors. In climates with a long resting period in winter, the plants will usually live much longer than in districts such as central and southern California where this period is confined to but one month of the year. Some may die after the first flowering, while others live for years.



PACIFIC STRAIN OF POLYANTHUS PRIMROSES

As the Polyanthus group is the most hardy and easiest to grow of all the Primrose family, succeeding in practically every climate, we have concentrated our work on improving this particular type. Our aim is to develop a wider and clearer color range, taller stems, and larger size of individual flowers and flower heads. After testing all of the important strains in commerce, we spent years of breeding and selecting from some 20,000 seedlings each year. We introduced our strain experimentally at the Golden Gate International Exposition on Treasure Island, where the exhibit, although comparatively small, created such interest that our stock was completely sold out within two weeks. This is, perhaps, the best recommendation our strain could have.

Selected Plants ... We can supply, from June to September, strong divisions of our breeding stock from the previous season, which are the most outstanding developments we have so far produced.

COLORS

White Orange Pastels Dark Red Golden Yellow Carmine-Rose Scarlet New Blends Bronze-Apricot

Price . . . 20c each; \$2.00 per dozen; \$12.00 per 100. F.O.B. Capitola.

Seedling Plants . . . During February and March we ship strong seedling plants, with flowers in assorted colors as per selected plants.

Price . . . \$1.50 per dozen; \$9.00 per 100. F.O.B. Capitola.

Seed . . . The seed we are offering is produced by artificial cross-pollination from the most outstanding new seedlings each year. By this method we can offer a quality which is in a class by itself. The production, naturally, is very limited and very costly but only in this way can the finest results be obtained. The new crop is available by July 1, which is also the best time for sowing. We pack approximately 125 seeds to a packet which, under reasonably good conditions, will germinate within two weeks after sowing. All colors listed as per selected plants are available separately or in complete mixture.

Price . . . \$1.00 per packet. (All seed sold out until July, 1942.)

THE CULTURE OF POLYANTHUS PRIMROSES

Seed... Seed of Polyanthus Primroses can be sown any time of the year, preferably in spring or early summer, so the seedling will reach flowering size by the next spring. Sow in any light mixture of soil; a good mixture is one-half leaf mold and one-half garden soil. Peat and garden soil with the addition of sand or any other combination can be

used so long as the soil is light and rich in humus. Cover the seed only slightly, moisten thoroughly, then cover the flats with burlap to keep them from drying out and place in heavy shade. When germination takes place, in about two weeks, give more light but protect from direct sunlight. Always keep moist. In six to eight weeks the young plants can be transplanted into flats again or directly into the open ground if large enough. In this case they should be protected from direct sunshine. The soil in the garden should be enriched with either old manure if sandy; or leaf mold, peat, and sand, if heavy. Additional feeding in early spring with any quick-acting fertilizer when the growth just commences. This should be preferably an organic fertilizer such as fishmeal or cottonseed meal which will help greatly to produce fine flowering specimens.

After two years the plants become too large and should be split and replanted in fresh soil. The best time for this is immediately after flowering. Plants transplanted late in the season will not get sufficient roots established in time for the flowering season and will then bring only poor flowers.

Primroses are hardy in every part of the United States. The only dangerous pest attacking them is the cutworm, of several varieties, which can be controlled with several baits sold commercially for that purpose.

VISIT OUR GARDENS

in March

when the Primroses are in bloom



CALLA LILIES

The cultural requirements of Calla Lilies are very simple. The pink and yellow Callas, being summer flowering plants, should be planted from January to April in good average garden soil, possibly in slightly shaded places in the warmer districts. Plant the bulbs approximately four inches deep and keep well supplied with moisture during the growing season and gradually withhold water after flowering, if possible. They can be left in the ground for years in regions with mild winters but should be lifted, dried, and stored in a frost-less place where winters are severe.

CALLA RHEMANII SUPERBA—New

The finest form of the pink Callas yet offered in cultivation. The seed of this species came from South Africa and it took a number of years to propagate it sufficiently for distribution. It is far superior to the so-called pink Callas that have so far been found in commerce. Flowers are fairly large, pale pink, plant very floriferous. For rock garden and cool borders this is a real acquisition. Delivery December to March.

Price . . . \$1.00 each; \$10.00 per dozen.

Calla Elliottiana . . . The yellow Calla Lily has larger flowers than the pink and is a very effective garden plant. Planting time February to April. Delivery of bulbs November to February.

Price . . . 25c each; \$2.50 per dozen; \$18.00 per hundred.



Ranunculus ... We have been interbreeding the Tecolote and the Australian strain which are the foremost of today, and are offering select stock resulting from these crosses. All single plants are weeded out, so that only uniform, selected plants are offered in the following color groups.

Tubers . . . White and Cream shades, Yellow and Gold shades, Apricot and Orange shades, Scarlet and Dark Red shades, Blush, Pink and Rose shades. Complete mixture. Delivery, July to November.

Price

Large tubers ... \$1.00 per doz.; \$7.00 per 100 Medium tubers ... 75c per doz.; \$5.00 per 100 Seed ... Ranunculus can readily be grown from seed, especially under California conditions. If sown during September or October and transplanted in December to permanent place, they will produce flowering specimens during March and April. In colder regions, only spring sowing can be recommended. PRICE ... 50c per packet.

Anemones (St. Brigid—Anderson Strain) . . . We have the pleasure of distributing the above-named strain, which is a distinct improvement on the types offered generally in commerce—not only in wider range of color but also much larger flowers, carried on stiff, strong stems. In warm regions Anemones can be had in bloom from November to end of March by planting from July to October. In colder regions early spring planting is prefer-

Price of Complete Mixture

Large tubers ... \$1.00 per doz.; \$7.00 per 100 Medium tubers ... 75c per doz.; \$5.00 per 100 Orders of six tubers supplied at dozen rate. Orders of 50 tubers supplied at 100 rate.

Seed . . . Anemones can be grown easily from seed and will bloom four to five months after sowing. Sowing time in California, from May to October. In regions with cold winters, early spring only.

PRICE . . . 50c per packet.

PLEASE USE THIS ORDER FORM

VETTERLE & REINELT

Hybridizing Gardens

CAPITOLA · CALIFORNIA

(Do not write on above line)

_By.

Received_

QUA

NAME_	(Please	print plainly)	ount Enclosed \$	
Addres	SS			
	ffice if different from mailing addres	ss		
	KINDLY LIST TUBERS, PLA	NTS AND	SEEDS SEPARATELY	7
YTITY	KIND OF TUBERS		COLOR	PRICE
California	Customers places and 3°/ Sales Tay			
Please do TERMS: C credit will CARRIAGI SUBSTITU	Customers please add 3% Sales Tarnot make remittance by stamps. Cash with order or approved refere be sent C. O. D. E: Please read carefully directions or TIONS: If sold out of any variety overwise specified on your order.	nces. Orders	de.	

PLANTS

COLOR

PRICE

KIND OF PLANTS

	SEI	DS	
QUANTITY	S E I	COLOR	PRICE
QUANTITY			PRICE

CARRIAGE

SEEDS... Are sent prepaid.

QUANTITY

BULBS... Add 25c packing and postage on all orders *under* \$2.50. If bulbs are sent at different times packing and postage charge is payable on each shipment of less than \$2.50.

PLANTS... Are shipped by *Express*. Carriage charge payable by purchaser. On orders of less than \$1.50 a packing charge of 25c must be added.

Please print or write your orders plainly and if writing us use separate sheet of paper.

PLEASE NOTE: Begonia Tubcrs available January to March. Begonia Seedlings available end May-June. Begonia Seeds available December 1st. Montbretia Tubers available December to March. Delphinium Plants available March-June and Sept.-November. Delphinium Seed available fresh crop July. Primrose Plants, Specimen Divisions available June to September. Primrose Seedlings available February and March. Primrose Seed available fresh crop July. Limited supply.

Tubers and Plants are charged at unit prices if labelled separately as to color etc.

If ordered at dozen rate will be shipped assorted.

In the event of any enquiry concerning your order, kindly give fullest particulars possible, as with our hundreds of orders this will help us and expedite replies. All stock guaranteed true to description but no guarantee for the result of growing over which we have no control.

PLEASE USE THIS ORDER FORM

VETTERLE & REINELT

Hybridizing Gardens

CAPITOLA · CALIFORNIA

(Do not write on above line)

By_

Filled_

Received

QUA

Date	(Please p	Am rint plainly)	ount Enclosed \$	
NAME_				
Addres	S			
Express offi	ice if different from mailing address			
	KINDLY LIST TUBERS, PLAN	NTS AND	SEEDS SEPARATE	LY
NTITY	KIND OF TUBERS		COLOR	PRICE
		-		
		-		
				-
	Customers please add 3% Sales Tax. not make remittance by stamps.		Amount Forward	
TERMS: Ca	ash with order or approved reference sent C. O. D.	ces. Orders	received without cash	or established
CARRIAGE	: Please read carefully directions or			
	FIONS: If sold out of any variety or wise specified on your order.	dered we sh	all supply a similar one YES	of equal value NO

PLANTS

COLOR

KIND OF PLANTS

PRICE

		, !	
	SEI	DS	
QUANTITY	KIND OF SEEDS	COLOR	PRICE
QUANTITY			PRICE
QUANTITY		COLOR	PRICE
QUANTITY	KIND OF SEEDS	COLOR	PRICE
QUANTITY	KIND OF SEEDS	COLOR	PRICE
QUANTITY	KIND OF SEEDS	COLOR	PRICE
QUANTITY	KIND OF SEEDS	COLOR	PRICE

CARRIAGE

SEEDS...Are sent prepaid.

OUANTITY

BULBS... Add 25c packing and postage on all orders *under* \$2.50. If bulbs are sent at different times packing and postage charge is payable on each shipment of less than \$2.50.

PLANTS... Are shipped by *Express*. Carriage charge payable by purchaser. On orders of less than \$1.50 a *packing* charge of 25c must be added.

Please print or write your orders plainly and if writing us use separate sheet of paper.

PLEASE NOTE: Begonia Tubers available January to March. Begonia Seedlings available end May-June. Begonia Seeds available December 1st. Montbretia Tubers available December to March. Delphinium Plants available March-June and Sept.-November. Delphinium Seed available fresh crop July. Primrose Plants, Specimen Divisions available June to September. Primrose Seedlings available February and March. Primrose Seed available fresh crop July. Limited supply.

Tubers and Plants are charged at unit prices if labelled separately as to color etc.

If ordered at dozen rate will be shipped assorted.

In the event of any enquiry concerning your order, kindly give fullest particulars possible, as with our hundreds of orders this will help us and expedite replies. All stock guaranteed true to description but no guarantee for the result of growing over which we have no control.

CULTURAL DIRECTIONS

RANUNCULUS

Seed... Seed can be sown from August to November in a mixture of light sandy loam. Cover the seed very lightly and keep moist and heavily shaded. When germination takes place, in from 30 to 60 days, give more light but protect from direct sunshine. Transplant when the second leaves are developed, either into flats, or if large enough, directly into open ground.

Location . . . Eastern exposure, with possible protection against hot afternoon sun, is best.

Soil . . . A rich, sandy loam, with perfect drainage.

Feeding ... Plant 4 to 6 inches apart and in one month, when plants are established, start feeding with quick-acting fertilizers—such as ammonium phosphate or fish meal. Rather small applications at intervals of a few weeks, instead of all at one time, are best.

Growing . . . Keep constantly moist, as they dislike drying out. After flowering, gradually withhold water and let the plants dry up. Dig and store the tubers in a cool, dry place. According to the time that flowers are wanted, plant tubers in October for January-February bloom: November-December for March-April flowering. In localities with severe winters, growing from seed is not practicable, as the young seedlings would freeze outdoors. Tubers stored in a cool, dry place can be planted very early in spring and bring satisfactory blooms in early summer. The cardinal points in cultivation are perfectly drained soil, constant moisture during growth, and protection from too-strong sunlight. Ranunculus are gross feeders, and to develop strong show plants one should avoid manures which help in keeping high moisture content in the soil, especially during rainy weather. They respond better to artificial feeding, as described above.

Pests . . . Birds like the young growth as it appears on the surface and it is advisable to protect the plants during the young stage with wire netting.

ANEMONES

Seed... Seed can be sown practically any time of the year in regions with mild winters, the sowing time being determined by the blooming season desired. Sow in a mixture of sandy leaf mold. Transplant when the second leaf develops, either in flats or in open ground. Use a light mixture of soil.

Location . . . Plants should have sufficient light, but never direct sunshine, for best results. The north side of the house is ideal.

Soil . . . Soil can be enriched heavily with well-rotted manure, as the plants can stand considerable moisture.

Growing . . . After flowering, gradual drying-off will ripen the tubers, which can be dug when all growth is withered. Stored in a cool, dry place, they will live for two years and can be planted when one wishes to have them in flower. The approximate time between planting and flowering is four months. In regions with severe winters, only spring planting is practicable or if planted in autumn they must be heavily covered to protect them from freezing. The natural flowering season is very early spring but in mild climates they can be had in bloom from November to the end of May. Anemones would not be practical to plant for summer flowering, as they dislike hot weather and the flowers would not develop properly.

Pests... Birds are again the most serious pest, eating young foliage. Wire netting will satisfactorily protect the young plants.

We have tried to make our growing suggestions as complete as possible.

To follow them exactly is not enough. Certain amount of knowledge must be gained by actual experience, and the use of common sense.



ARRANGEMENT OF TUBEROUS BEGONIAS